INDEX OF AUTHORS AND SUBJECTS

VOL. 84

Compiled by A.R. McGILL

AHLQUIST, J.E., see C.G. SIBLEY
Albatross, Wandering, Some aspects of the morphology on Macquarie Island, 29–32
AMBROSE, S.J., The response of small birds to extreme heat, 242–243
Babbler, Grey-crowned, measuring growth in nestlings, 185–187; Communal building of brood and roost nests, 193–199
BAIRD, R.F., The Pleistocene distribution of the Tasmanian Native-hen *Gallinula mortierii mortierii*, 119–123
BELL, H.L., A bird community of lowland rainforest in New Guinea, 6. Foraging ecology and community structure of the avifauna, 142–158; Polygamy in the Speckled Warbler *Sericornis sagittatus*, 183–184; The importance of foothill forest in the diversity of rainforest birds in New Guinea, 225–235; A note on communal breeding and dispersal of young of the Hooded Robin *Petroica cucullata*, 243–244
Biology of the Little Eagle on the Northern Tablelands of N.S.W., 87–92
Bird community of lowland rainforest in New Guinea, 6. Foraging ecology and community structure of the avifauna, 142–158
BOOTH, D.T. and R.S. SEYMOUR, Effect of adding water to Malleefowl mounds during a drought, 116–118
Breeding biology of the Australasian Gannet *Morus serrator* (Gray) at Motu Karamarama, Hauraki Gulf, N.Z., 1. The egg, 129–136; II. Breeding success and chick growth, 211–224
BRIFFA, P., see B.G. COLLINS
CAMPBELL, N.A., see D.A. SAUNDERS
CANNON, C.E., The diet of lorikeets *Trichoglossus* spp in the Queensland-New South Wales border region, 16–22
Citation, R.A.O.U. Fellow, S.G. Lane, 124
CLARKE, M.F., Interspecific aggression within the genus *Manorina*, 113–115
CLOSE, D. and O.M.G. NEWMAN, The decline of the Eastern Curlew in south-eastern Australia, 38–40
COLLINS, B.G., P. BRIFFA and C. NEWLAND, Temporal changes in abundance and resource utilization by honeyeaters at Wongamine Nature Reserve, 159–166
Communal building of brood and roost nests by the Grey-crowned Babbler, 193–199
Coucal, Pheasant, Geographical variation and a description of a new subspecies from Timor, 1–15
Curlew, Eastern, the decline in south-eastern Australia, 38–40
CURRY, P.J., see D.L. SERVENTY
DAWSON, T.J., D. READ, E.M. RUSSELL and R.M. HERD, Seasonal variation in daily activity patterns, water relations and diet of Emus, 93–102
DEBUS, S., Biology of the Little Eagle on the Northern Tablelands of New South Wales, 87–92
DENNISON, M.D., see H.A. ROBERTSON
De REBEIRA, C.P., see R.D. WOOLLER
Diet of lorikeets *Trichoglossus* spp in the Queensland-N.S.W. border region, 16–22
DOW, D.D. and B.J. GILL, Measuring growth in nestling Grey-crowned Babblers, 185–187
DOW, D.D. and B.R. KING, Communal building of brood and roost nests by the Grey-crowned Babbler *Pomatostomus temporalis*, 193–199
Duck, Pacific Black, heavy metal residues in, from Griffith, N.S.W., 44–46
  kleptoparasitism, including size-related differences, 65–70
DUNN, K., see T. STOKES
DUDZINSKI, M.L., see I.J. MASON
Eagle, Little, biology, on the Northern Tablelands of N.S.W., 87–92
Egg shape within the Australian Psittaciformes with comments on eggs of *Nymphicus hollandicus*, 36–37
Emu, Seasonal variations in daily activity patterns, water relations and diet, 93–102
Fairy-wren, Campbell's five specimens from New Guinea, 249–250
  Superb, territory quality and group size, 178–180
Forest avifauna near Bega, N.S.W., I. Differences between forest types, 200–210
FUJIMAKI, Y. and I.J. SKIRA, Notes on Latham's Snipe *Gallinago hardwickii* in Japan, 49–51
Gannet, Australasian, breeding biology at Motu Karamarama, Hauraki Gulf, N.Z., I. The egg, 129–136; II. Breeding success and chick growth, 211–224
Geographical variation in the Pheasant Coucal and a description of a new subspecies from Timor, 1–15
GILL, B.J., see D.D. DOW
Hardhead, kleptoparasitism, including size-related differences, 65–70
HERD, R.M., see T.J. DAWSON
Honeyeater, White-plumed, protection behaviours of breeding, 42–43
Honeyeaters, small birds, lerp-feeding and the problem of, 137–141
  temporal changes in abundance and resource utilization, 159–166
HURLEY, V.G., see F.I. NORMAN
IMBER, M.J., Migration of White-faced Storm-Petrels *Pelagodroma marina* in the South Pacific and the status of the Kermadec subspecies, 32–35
Importance of foothill forest in the diversity of rainforest birds in New Guinea, 225–235
Interbreeding of the Spotted and Yellow-rumped Pardalotes *Pardalotus punctatus* and *P. xanthopygus*, 80–86
International Commission on Zoological Nomenclature, 62–63, 256
JOUVENTIN, P. and J.P. ROBIN, Olfactory experiments on some Antarctic birds, 46–48
KING, B.R., The Herald Petrel *Pterodroma arminjoniana heraldica* breeding on Raine Island, Qld., 246–247
  see D.D. DOW
Kleptoparasitism in Hardheads and Pacific Black Ducks, including size-related differences, 65–70
Lorikeets, *Trichoglossus* spp, diet in the Qld-N.S.W. border region, 16–20
McFARLAND, D., Protection behaviours of breeding White-plumed Honeyeaters *Lichenostomus penicillatus*, 42–43
McKEAN, J.L., see I.J. MASON
Malleefowl, effect of adding water to mounds during a drought, 116–118
Manorina, interspecific aggression within the genus, 113–115
MARCHANT, S., Nest records of the Eastern Yellow Robin *Eopsaltria australis*, 167–174
MASON, I.J., J.L. McKEAN and M.L. DUDZINSKI, Geographical variation in the Pheasant Coucal *Centropus phasianinus* (Latham) and a description of a new subspecies from Timor, 1–15
MONTAGUE, T.L., The food of Antarctic Petrels *Thalassoica antarctica*, 244–245
Native-hen, Tasmanian, the Pleistocene distribution, 119–123

NEWLAND, C., see B.G. COLLINS
NEWMAN, O.M.G., see D. CLOSE
NIAS, R.C., Territory quality and group size in the Superb Fairy-wren *Malurus cyaneus*, 178–180
NORMAN, F.I. and V.G. HURLEY, Gonad measurements and other parameters from Chestnut Teal *Anas castanea* collected in the Gippsland Lakes region, Victoria, 52–55
Nymphicus hollandicus, comments on the eggs, 36–37
Olfactory experiments on some Antarctic birds, 46–48
OLSEN, P.D., Heavy metal residues in Pacific Black Duck *Anas superciliosa* from Griffith, New South Wales, 44–46
Pardalotes, Spotted and Yellow-rumped, interbreeding, 80–86
PARKER, S.A., The identity of *Sericornis tyrannula* de Vis, 108–110
Peltops, Papuan genus, the relationships, 181–183
Petrel, Antarctic, the food, 244–245
Herald, breeding on Raine Island, Qld., 246–247
READ, D., see T.J. DAWSON
Relationships of the Australo-Papuan Treecreepers Climacteridae as indicated by DNA-DNA hybridization, 236–241
of the Papuan genus *Peltops*, 181–183
Response of small birds to extreme heat, 242–243
REVIEWS, 56–62, 125–127, 188–190, 251–256
ROBERTSON, H.A. and M.D. DENNISON, Sexual dimorphism of the Chatham Island Warbler *Gerygone albofrontata*, 103–107
ROBIN, J.P., see P. JOUVENTIN
Robin, Eastern Yellow, nest-records, 167–174
Hooded, a note on communal breeding and dispersal of young, 243–244
RUSSELL, E.M., see T.J. DAWSON
SAUNDERS, D.A., G.T. SMITH and N.A. CAMPBELL, Egg shape within the Australian Psittaciformes with comments on eggs of *Nymphicus hollandicus*, 36–37
SAUNDERS, D.A., see R.D. WOOLLER
SCHODDE, R., First specimens of Campbell's Fairy-Wren *Malurus campbelli* from New Guinea, 249–250
SCHODDE, R., see C.G. SIBLEY
Seasonal variation in daily activity patterns, water relations and diet of Emus, 93–102
Seed-eating birds, a non-destructive method to determine the diet, 40–41
Sericornis tyrannula de Vis, the identity, 108–110
SERVENTY, D.L. and P.J. CURRY, Observations on colony size, breeding success, recruitment and inter-colony dispersal in a Tasmanian colony of Short-tailed Shearwaters *Puffinus tenuirostris* over a 30-year period, 71–79
Sexual dimorphism of the Chatham Island Warbler *Gerygone albofrontata*, 103–107
SEYMOUR, R.S., see D.T. BOOTH
Shearwater, Short-tailed, observations on colony size, breeding success, recruitment and inter-colony dispersal in a Tasmanian colony over a 30-year period, 71–79
SHEILS, W., see T. STOKES
Shelduck, Australian, consistent individuality in the calls on Rottnest Is., W.A. 175–177
SIBLEY, C.G. and J.E. AHLQUIST, The relationships of the Papuan genus *Peltops*, 181–183
SIBLEY, C.G., R. SCHODDE and J.E. AHLQUIST, The relationships of the Australo-Papuan Treecreepers Climacteridae as indicated by DNA-DNA hybridization, 236–241
SKIRA, I.J., Breeding distribution of the Brown Skua on Macquarie Island, 248–249
SKIRA, I.J., see Y. FUJIMAKI
Skua, Brown, breeding distribution on Macquarie Island, 248–249
Small birds, lerp-feeding and the problem of honeyeaters, 137–141
SMITH, G.T., see D.A. SAUNDERS
SMITH, P., The forest avifauna near Bega, New South Wales, I. Differences between forest types, 200–210
Snipe, Latham's, notes in Japan, 49–51
STOKES, T., W. SHEILS and K. DUNN, Birds of the Cocos (Keeling) Islands, Indian Ocean, 23–28
Storm-Petrel, White-faced, migration in the South Pacific and the status of the Kermadec subspecies, 32–35
STRAW, B., see R. ZANN
Swamphen, Purple, simultaneous polyandry, 111–113
Teal, Chestnut, gonad measurements and other parameters in the Gippsland Lakes region, Vic., 52–55
TOMKINS, R.J., Some aspects of the morphology of Wandering Albatrosses on Macquarie Island, 29–32
Treecreepers Climacteridae, the relationships as indicated by DNA-DNA hybridization, 236–241
Warbler, Chatham Island, sexual dimorphism, 103–107
Speckled, polygamy, 183–184
WETTIN, P., Simultaneous polyandry in the Purple Swamphen, 111–113
WINGHAM, E.J., Breeding biology of the Australasian Gannet *Morus serrator* (Gray) at Motu Karamarama, Hauraki Gulf, New Zealand I. The egg, 129–136; II. Breeding success and chick growth, 211–224
WOINARSKI, J.C.Z., Interbreeding of the Spotted and Yellow-rumped Pardalotes *Pardalotus punctatus* and *P. xanthopygus*, 80–86; Small birds, lerp-feeding and the problem of honeyeaters, 137–141
WOODALL, P.F., Kleptoparasitism in Hardheads and Pacific Black Ducks, including size-related differences, 65–70
WOOLLER, R.D., D.A. SAUNDERS and C.P. De REBEIRA, Consistent individuality in the calls of Australian Shelducks *Tadorna tadornoides* on Rottnest Island, Western Australia, 175–177
ZANN, R. and B. STRAW, A non-destructive method to determine the diet of seed-eating birds, 40–41

INDEX TO SCIENTIFIC NAMES

VOL. 84

Compiled by A.R. McGILL

Acanthiza, 173
—— *apicalis*, 137–140
—— *chrysorrhoa*, 181, 198, 238–240, 243
—— *inornata*, 238–239
—— *lineata*, 137–140, 209
—— *nana*, 137–140
—— *pusilla*, 115, 137–140, 209
—— *reguloides*, 137–140, 209, 243
—— *uropygialis*, 137–140
Acanthorhynchus superciliosus, 159, 239
—— *tenuirostris*, 209
Accipiter buergersi, 232
—— *cirrhocephalus*, 206, 232
—— *fasciatus*, 14, 88, 120, 206, 232
—— *novaehollandiae*, 206, 232
—— *poliocephalus*, 232
Aegotheles bennettii, 233
—— *cristatus*, 207
Aepypodius arfakianus, 232
Ailuroedus buccoides, 146, 235, 238–239
—— *crassirostris*, 238–239
Alcedo atthis, 229
—— *azurea*, 233
—— *pusilla*, 233
Alisterus chloropterus, 147, 233
—— *scapularis*, 207
Amaurornis olivaceus, 232
Amazona vittata, 253
Amytornis striatus, 250
Anas americana, 65
—— *castanea*, 52–55
—— *erythrorhyncha*, 53
—— *gibberifrons*, 88
—— *penelope*, 65
—— *platyrhynchos*, 44
—— *strepera*, 65
—— *superciliosa*, 44–46, 65–70
—— *waigiuensis*, 229
Anhinga melanogaster, 232
Anous stolidus, 27
Anseranas semipalmata, 198
Anthochaera carunculata, 115, 139, 209, 238–240
—— *chrysoptera*, 139, 209, 239
Aplonis cantoroides, 235
—— *metallica*, 146, 235
Aprosmictus, 36
—— *erythropterus*, 14
Apus pacificus, 207
Aquila audax, 90, 206
Ardea alba, 232
—— *novaehollandiae*, 25, 206
Ardeola ibis, 25
Arenaria interpres, 26
Arses telescophthalmus, 144, 234
Artamus cinereus, 14, 181, 238
—— *cyanopterus*, 181, 210
—— *leucorhynchus*, 235
—— *superciliosus*, 210, 238–240
Ashbyia lovensis, 239
Atrichornis clamosus, 59
Aviceda cuculoides, 252
—— *subcristata*, 232
Aythya australis, 65–70, 188
Barnardius, 36
Buteo galapagoensis, 126
Cacatua galerita, 14, 233
—— *roseicapilla*, 36
Cacomantis castaneiventris, 233
—— *variolosus*, 8, 233
Calidris acuminata, 56
—— *alba*, 26
—— *ruficollis*, 56
Caliechthrus leucolophus, 233
Callocephalon, 36
—— *fimbriatum*, 207
Calyptorhynchus funereus, 36, 207
—— *lathami*, 207
Campephaga quiscalina, 181
Campochaera sloetii, 147, 233
Campylorhynchus, 193
Caprimulgus macrourus, 233
—— *mystacalis*, 207
Cardellina, 188
Carduelis carduelis, 209
Casuarius bennetti, 232
—— *casuarius*, 232
Cecropis ariel, 208
—— *nigricans*, 208
Centropus ateralbus, 11–13
—— *bengalensis*, 12–13
—— *bernsteini*, 11–13
—— *chalybeus*, 11
—— *goliath*, 11–12
—— *menbeki*, 11, 233
—— *milo*, 11–12
—— *phasianinus*, 1–15, 233
—— *spilopterus*, 2–13
—— *toulou*, 11
—— *violaceus*, 11
Certhia familiaris, 238–240
Certhionyx variegatus, 238
Ceyx azureus, 207
—— *lepidus*, 233
Chaetorhynchus papuensis, 235
Chalcophaps indica, 147, 232
—— *stephani*, 147, 232
Chalcopsitta scintillata, 147, 232
Charmosyna placentis, 147, 232
—— *pulchella*, 155, 232
Chenonetta jubata, 88
Chettusia leucura, 56
Chlamydera cerviniventris, 235, 238
—— *lauterbachi*, 238
—— *maculata*, 238–239
—— *nuchalis*, 181, 238–240
Chlidonias leucoptera, 26
Chrysococcyx, 144
—— *lucidus*, 207
—— *meyerii*, 233
—— *minutillus*, 8, 233
—— *osculans*, 183
Cicinnurus regius, 146, 235
Cinclosoma ajax, 233
—— *cinnamomeum*, 181, 238–239
—— *punctatum*, 208
Circus aeruginosus, 25, 120, 206
—— *assimilis*, 120
Cisticola exilis, 234
Climacteris affinis, 236–241
—— *erythrops*, 208, 236–241
—— *leucophaea*, 174, 208
—— *melanura*, 236–241
—— *picumnus*, 236–241, 243
—— *rufa*, 236–241
Clytoceyx rex, 233
Collocalia esculenta, 144, 233
—— *vanikorensis*, 144, 233
Colluricincla harmonica, 115, 208, 238
—— *megarhyncha*, 144, 234
Columba livia, 57
—— *vitiensis*, 232
Conopophila whitei, 239
Coracina boyeri, 147, 233
—— *caeruleogrisea*, 233
—— *caesia*, 181
—— *lineata*, 233
—— *melaena*, 146, 233
—— *novaehollandiae*, 181, 208, 233
—— *papuensis*, 181, 208, 233
—— *schisticeps*, 233
—— *tenuirostris*, 208, 233
Corcorax melanorhamphus, 199, 238–240
Cormobates leucophaea, 236–241
—— *placens*, 236–241
Corvinella corvina, 181
Corvus corax, 69
—— *coronoides*, 88, 210
—— *mellori*, 239–240
—— *orru*, 235
—— *tristis*, 235
Coturnix australis, 206
—— *ypsilophorus*, 14
Cracticus cassicus, 235
—— *quoyi*, 181, 235
—— *torquatus*, 210
Crateroscelis gutturalis, 230
—— *murina*, 234
Cuculus pallidus, 207
—— *pyrrhophanus*, 207
—— *saturatus*, 233
—— *variolosus*, 207
Dacelo gaudichaud, 233
—— *gigas*, 198
—— *novaeguineae*, 171, 207
Daphoenositta chrysoptera, 115, 209, 238–240
Daption capense, 46, 245
Dendroica, 253
Dicaeum hirundinaceum, 209
—— *pectorale*, 234
Dicrurus adsimilis, 181
—— *hottentottus*, 146, 209, 235
—— *paradiseus*, 181, 238
Diomedea exulans, 29–32
Diphyllodes magnificus, 235
Domicella lory, 147
Dromaius novaehollandiae, 88, 93–102, 168

Drymodes brunneopygia, 238–240
—— *superciliaris*, 234
Ducula bicolor, 232
—— *mullerii*, 147, 232
—— *pinon*, 232
—— *rufigaster*, 147, 232
—— *whartoni*, 27
—— *zoeae*, 147, 232
Eclectus, 36
—— *roratus*, 233
Egretta sacra, 25
Elanus notatus, 206
Emblema oculata, 41
—— *temporalis*, 115, 209
Eopsaltria australis, 115, 167–174, 208
—— *griseogularis*, 239–240
Ephthianura albifrons, 239
—— *tricolor*, 242
Epimachus albertisi, 235
Ergaticus, 188
Erithacus rubecula, 173
Erythrura trichroa, 235
Eudynamys scolopacea, 8, 147, 233
Eudyptes chrysolophus, 248
Eudyptula, 188
Eulabeornis plumbeiventris, 232
Eupetes caerulescens, 144
Eurostopodus papuensis, 233
Eurynorhynchus pygmaeus, 56
Eurystomus orientalis, 144, 207, 233
Euthlypis, 188
Falco, 59
—— *berigora*, 206
—— *cenchroides*, 206
—— *longipennis*, 206
—— *peregrinus*, 206, 232
—— *severus*, 232
—— *sparverius*, 126
Falcunculus frontatus, 115, 208
Fratercula arctica, 69
Fregata ariel, 24, 246
—— *minor*, 24, 246
Fulica atra, 66
Fulmarus glacialis, 48, 77
—— *glacialoides*, 46, 245
Gallicolumba jobiensis, 232
—— *rufigula*, 147, 232
Gallinago gallinago, 26, 50
—— *hardwickii*, 49–51
—— *stenura*, 26
Gallinula mortierii, 111, 119–123
Gallirallus australis, 249
Gallus gallus, 25
Gavia pacifica, 56
Geoffroyus geoffroyi, 147, 233
—— *simplex*, 155, 233
Geopelia placida, 207
—— *striata*, 14, 57
Geophaps scripta, 57
Gerygone, 173
—— *albofrontata*, 103–107
—— *chloronota*, 234
—— *chrysogaster*, 144, 234
—— *fusca*, 103–105, 137, 239
—— *igata*, 103
—— *laevigaster*, 105
—— *magnirostris*, 234
—— *mouki*, 106, 137–140, 209
—— *olivacea*, 137, 209
—— *palpebrosa*, 105, 150, 234
Glareola maldivarum, 26
Glossopsitta, 36
—— *concinna*, 207
—— *pusilla*, 207
Glycichaera fallax, 234
Goura scheepmakeri, 232
Grallina bruijni, 235
—— *cyanoleuca*, 14, 168, 181, 210, 238–240, 242
Granatellus, 188
Gygis alba, 27
Gymnocrex plumbeiventris, 144
Gymnophaps albertisii, 147, 232
Gymnorhina tibicen, 88, 181, 210, 238
Halcyon macleayii, 233
—— *sancta*, 207, 233
—— *torotoro*, 233
Haliaeetus leucogaster, 206, 232
Haliastur indus, 232
—— *sphenurus*, 206, 232
Halobaena caerulea, 249
Harpyopsis novaeguineae, 232
Hemiprocne mystacea, 144, 233
Henicopernis longicauda, 232
Henicophaps albifrons, 232
Hieraaetus dubius, 91
—— *fasciatus*, 91
—— *morphnoides*, 87–92, 206, 232
—— *pennatus*, 91
—— *wahlbergi*, 90
Hirundapus caudacutus, 27, 207, 233
Hirundo neoxena, 167, 207
—— *rustica*, 27, 173
—— *tahitica*, 233
Ibidorhyncha struthersii, 56
Icteria, 188
Jacana spinosa, 111
Lagonosticta senegala, 40
Lagopus scoticus, 43
Lalage leucomela, 146, 233
—— *nigra*, 181
—— *sueurii*, 14, 181
Laniarius barbarus, 181
Lanius excubitor, 181
Larius roratus, 147
Larus, 69
Lathamus, 36
Leipoa ocellata, 116–118
Leucopeza, 188
Leucosarcia melanoleuca, 207
Lichenostomus chrysops, 115, 139, 209
—— *fuscus*, 43, 139, 209
—— *leucotis*, 43, 139, 209
—— *melanops*, 43, 139, 209
—— *obscurus*, 234
—— *ornatus*, 139
—— *penicillatus*, 42–43, 115, 139
—— *virescens*, 159
Lichmera indistincta, 14, 159–165, 239
Limicola falcinellus, 56
Limnodromus scolopaceus, 56
Limnothlypis, 188
Lonchura caniceps, 235
—— *tristissima*, 235
Lophoictinia isura, 206
Lopholaimus antarcticus, 206
Loriculus aurantiifrons, 233
Lorius lory, 232
Machaerirhynchus flaviventer, 144, 234
Macheiramphus alcinus, 232
Macronectes giganteus, 46
—— *halli*, 60
Macropygia, 147
—— *amboinensis*, 145, 206, 232
—— *nigrirostris*, 232
—— *phasianella*, 57
Malaconotus cruentus, 181
Malurus alboscapulatus, 234
—— *campbelli*, 249–250
—— *cyaneus*, 89, 115, 178–180, 193, 208
—— *cyanocephalus*, 234
—— *grayi*, 249–250
—— *lamberti*, 208
—— *leucopterus*, 242
—— *splendens*, 198, 238–239, 242
Manorina melanocephala, 113–115, 137–139, 193
—— *melanophrys*, 113–115, 198, 209
Manucodia ater, 146, 235
—— *chalybata*, 235
—— *keraudrenii*, 146, 235
Mearnsia novaeguineae, 233
Megacephalon maleo, 228
Megapodius freycinet, 232
Megatriorchis doriae, 232
Melanocharis nigra, 146, 234
—— *versteri*, 181, 238
Melidora macrorrhina, 233
Melilestes megarhynchus, 146, 234
Meliphaga, 146
—— *albonotata*, 234
—— *analoga*, 234
—— *aruensis*, 234, 240
—— *gracilis*, 234
—— *lewinii*, 209, 238
—— *orientalis*, 234
—— *penicillata*, 239
Melithreptus affinis, 140
—— *brevirostris*, 139, 164, 209
—— *lunatus*, 115, 139, 209
Melopsittacus, 36
—— *undulatus*, 242
Menura novaehollandiae, 207, 238–239
Merops ornatus, 144, 233, 242
Microdynamis parva, 147, 233
Microeca brunneicauda, 110
—— *griseoceps*, 234
—— *flavovirescens*, 144, 234
—— *leucophaea*, 208
Microligea, 188
Micropsitta bruijnii, 233
—— *pusio*, 146, 233
Mino anais, 147, 235
—— *dumonti*, 147, 235
Monachella muelleriana, 234
Monarcha chrysomela, 144, 234
—— *cinerascens*, 181
—— *frater*, 234
—— *guttula*, 144, 234
—— *melanopsis*, 144, 181, 208, 234
—— *trivirgata*, 156
Morus bassana, 129–135, 211
—— *capensis*, 129–135, 211
—— *serrator*, 129–136, 211–224
Motacilla, 27
Myiagra alecto, 144, 181, 234, 238–239
—— *cyanoleuca*, 208
—— *inquieta*, 115, 208
—— *rubecula*, 208, 234
Myzomela adolphinae, 234
—— *cruentata*, 234
—— *eques*, 234

—— *nigrita*, 234
—— *obscura*, 239
—— *rosenbergii*, 234
—— *sanguinolenta*, 209
Nectarinia jugularis, 234
—— *sericea*, 146, 234
Neophema, 36
—— *chrysogaster*, 59
Nephelornis, 188
Ninox novaeseelandiae, 14, 207
—— *rufa*, 233
—— *strenua*, 207
—— *theomacha*, 233
Northiella, 36
Numenius madagascariensis, 38–40, 56
—— *minutus*, 26, 86
Numida meleagris, 25
Nycticorax caledonicus, 25
Nymphicus hollandicus, 36–37
Oceanites oceanicus, 46
Oceanodroma leucorhoa, 34, 46
Oedistoma iliolophum, 234
—— *pygmaeum*, 234
Opopsitta gulielmitertii, 147, 233
Oriolus chinensis, 181
—— *cruentus*, 181
—— *sagittatus*, 181, 209
—— *szalayi*, 146, 235
Orthonyx spaldingi, 230
—— *temminckii*, 238–240
Otidiphaps nobilis, 232
Pachycare flavogrisea, 234
Pachycephala, 174
—— *aurea*, 228
—— *hyperythra*, 234
—— *leucogaster*, 234
—— *olivacea*, 208
—— *pectoralis*, 115, 208, 238–240
—— *rufiventris*, 115, 181, 208, 239
—— *simplex*, 144, 234
Pachycephalopsis poliosoma, 234
Pachyptila desolata, 249
—— *turtur*, 249
Padda, 58
—— *oryzivora*, 27
Pagodroma nivea, 46, 245
Parabuteo unicinctus, 126
Paradisaea minor, 238
—— *raggiana*, 146, 235
Pardalotus punctatus, 80–86, 115, 137–140, 209
—— *quadragintus*, 140
—— *striatus*, 137–140, 209, 238–240
—— *xanthopygus*, 80–86, 137
Parmoptila, 58
Passer domesticus, 69
Pelagodroma marina, 32–35
Pelecanoides urinatrix, 249
Peltops blainvillii, 144, 181, 234
—— *montanus*, 181–183
Pericrocotus cinnamomeus, 181
Petrochelidon nigricans, 233
Petroica, 106
—— *cucullata*, 239, 243–244
—— *multicolor*, 208
—— *phoenicea*, 208
—— *rosea*, 115, 208
Peucedramus, 188
Pezoporus, 36
Phaethon lepturus, 25
—— *rubricauda*, 24, 246
Phalacrocorax sulcirostris, 232
Phaps chalcoptera, 207
—— *elegans*, 207
Philemon buceroides, 235
—— *citreogularis*, 14
—— *corniculatus*, 209
—— *meyeri*, 235
Phylidonyris albifrons, 239
—— *nigra*, 43, 159–165, 203
—— *novaehollandiae*, 43, 115, 209
—— *pyrrhoptera*, 209
Phylloscopus trivirgatus, 234
—— *trochilus*, 56
Pitohui cristatus, 234
—— *dichrous*, 234
—— *ferrugineus*, 146, 181, 234
—— *kirhocephalus*, 238
Pitta erythrogaster, 144, 233
—— *sordida*, 144, 233
—— *versicolor*, 230
Pityriasis gymnocephala, 181
Platycercus, 36
—— *elegans*, 207
—— *eximius*, 115, 207
Pluvialis dominica, 26
Podargus ocellatus, 233
—— *papuensis*, 233
—— *strigoides*, 207
Poecilodryas hypoleuca, 144, 234
—— *placens*, 228
Poephila guttata, 14, 40, 242
Polytelis, 36
Pomatostomus halli, 60, 199
—— *isidorei*, 144, 234
—— *ruficeps*, 199
—— *superciliosus*, 199, 239
—— *temporalis*, 185–187, 193–199
Porphyrio porphyrio, 111–113
Prionops plumata, 181
Probosciger, 36
—— *aterrimus*, 233
Procellaria passerina, 34
Psephotus, 36
Pseudeos fuscata, 146, 232
Psitteuteles, 36
Psophodes olivaceus, 174, 208, 230
Pteridophora alberti, 181
Pterodroma arminjoniana, 246–247
—— *lessoni*, 249
Ptilinopus coronulatus, 147, 232
—— *dupetithouarsii*, 57
—— *iozonus*, 147
—— *magnificus*, 57, 147, 232
—— *nanus*, 147, 232
—— *ornatus*, 232
—— *perlatus*, 147, 232
—— *pulchellus*, 232
—— *regina*, 57
—— *rivoli*, 232
—— *superbus*, 147, 232
Ptilonorhynchus violaceus, 210, 238–240
Ptiloris magnificus, 146, 235, 238–239
Ptilorrhoa caerulescens, 233
—— *castanonota*, 234
Puffinus carneipes, 32
—— *gravis*, 46
—— *griseus*, 77, 249
—— *pacificus*, 24
—— *puffinus*, 77
—— *tenuirostris*, 71–79, 245
—— *paradiseus*, 240
Purpureicephalus, 36
Pycnopygius ixoides, 235
—— *stictocephalus*, 235
Pycnoptilus floccosus, 188, 208
Pygoscelis adeliae, 248
Rallina tricolor, 232
Rallus philippensis, 25
Ramsayornis modestus, 239
Raphus cucullatus, 60
Reinwardtoena reinwardtsii, 147, 232
Rhabdornis, 236
Rhamphomantis megarhynchus, 233
Rhipidura fuliginosa, 115, 138–139, 156, 208
—— *hyperythra*, 144, 234
—— *leucophrys*, 115, 167, 208, 234, 238–240, 242
—— *leucothorax*, 234
—— *maculipectus*, 234
—— *rufidorsa*, 144, 234
—— *rufifrons*, 181, 208
—— *rufiventris*, 144, 234
—— *threnothorax*, 144, 234
Rhodostethia rosea, 56
Rhyticeros plicatus, 233
Salpornis, 238
Scythrops novaehollandiae, 203, 233
Seleucidis melanoleuca, 235
Sericornis cautus, 108
—— *citreogularis*, 208
—— *frontalis*, 108, 115, 208, 238–240, 242
—— *keri*, 60
—— *laevigaster*, 108
—— *lathami*, 230
—— *magnirostris*, 208
—— *nouhuysi*, 239
—— *pyrrhopygius*, 108–110
—— *sagittatus*, 183–184
—— *spilodera*, 234
—— *tyrannulus*, 108–110
Sericulus aureus, 227
Setophaga, 188
Sipodotus wallacii, 234, 250
Sitta carolinensis, 238–240
Smierornis brevirostris, 137–140, 239
Somateria mollisima, 65
Sphecotheres viridis, 181
Stercorarius maccormicki, 47
—— *parasiticus*, 69
—— *skua*, 248–249
Sterna bergii, 44
—— *fuscata*, 26
Stipiturus, 250
Strepera graculina, 171, 181, 120
—— *versicolor*, 181, 210
Streptopelia chinensis, 57, 206
—— *senegalensis*, 57
Struthidea cinerea, 199, 238–240
Sturnus vulgaris, 88
Sula dactylatra, 24, 246
—— *leucogaster*, 24
—— *sula*, 24, 246
Syma torotoro, 144
Tadorna tadorna, 175
—— *tadornoides*, 175–177
Tallegalla fuscirostris, 232
Tanysiptera danae, 233
—— *galatea*, 144, 233
—— *sylvia*, 233
Tchagra australis, 181

Tephrodornis gularis, 181
—— *pondicerianus*, 181
Teretistris, 188
Thalassoica antarctica, 244–245
Timeliopsis griseigula, 146, 234
Todopsis cyanocephala, 144
Toxorhamphus iliolophus, 146
—— *poliopterus*, 234
Tregellasia leucops, 181, 234, 238
Tribonyx mortierii, 198
Trichoglossus, 36
—— *chlorolepidotus*, 16–22
—— *goldiei*, 232
—— *haematodus*, 16–22, 146, 207, 232
Tringa hypoleucos, 26, 232
—— *nebularia*, 26
Trugon terrestris, 147, 232
Turdus poliocephalus, 27
Turnix varia, 206
Tyto alba, 207, 233
—— *tenebricosa*, 207
Uraeginthus, 58
Uroglaux dimorpha, 233
Vireo griseus, 181
Xanthotis flaviventer, 146, 235
—— *polygramma*, 146, 235
Xenoligea, 188
Zoothera dauma, 115, 208, 230
Zosterops lateralis, 209
—— *natalis*, 27
—— *novaeguineae*, 234

ERRATA

Page 16 *Trichoglossus haematodotus* should be *T. haematodus*
16 *Trichoglossus chlorolepitodus* should be *T. chlorolepidotus*
46 *Oceanodroma leucorrhoa* should be *Oceanodroma leucorhoa*. It is spelt correctly on page 34
56 *Namenius* should be *Numenius*
105 *Gerygone palpebrata* should be *G. palpebrosa*
111 *Gallinula morterii* should be *G. mortierii* again spelt erroneously on page 111 where *T.* indicates *Tribonyx??*
126 *Parabuteo unicintus* should be *P. unicinctus*
126 *Buteo galapogoensis* should be *B. galapagoensis*
144 *Pomatostomus isidori* should be *P. isidorei*
146 *Pitohui ferruginea* should be *P. ferrugineus*
147 *Trugon terrestis* should be *T. terrestris*
207 *Ceyx azurea* should be *C. azureus*
232 *Reinwardtoena reinwardtii* should be *R. reinwardtsii*
233 *Hirundapus caudacuta* should be *Hirundapus caudacutus*
235 *Manucodia atra* should be *M. ater*